

# Anti-uracil-DNA glycosylase antibody



Catalog Number: 101035

## Product name

Anti-uracil-DNA glycosylase antibody

IHC-P: 0.1-2 µg/mL

ICC/IF: 0.5-1.5 µg/mL

## Specificity

Human uracil-DNA glycosylase / UNG

## Validations

## Antibody description

Rabbit polyclonal to uracil-DNA glycosylase

## Preparation

Produced in rabbits immunized with a synthetic peptide corresponding to the center region of the Human uracil-DNA glycosylase / UNG, and purified by antigen affinity chromatography.

## Formulation

0.2 µm filtered solution in PBS

## Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C.

Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

## Clonality

Polyclonal


## Ig Type

Rabbit IgG

## Applications


IHC-P, IF, ICC/IF

## Dilutions



uracil-DNA glycosylase / UNG Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Immunochemical staining of human UNG in human stomach with rabbit polyclonal antibody (0.5 µg/mL, formalin-fixed paraffin embedded



sections).

uracil-DNA glycosylase / UNG Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Immunochemical staining of human UNG in human testis with rabbit polyclonal antibody (0.5 µg/mL, formalin-fixed paraffin embedded

# Anti-uracil-DNA glycosylase antibody




Catalog Number: 101035

---

uracil-DNA glycosylase / UNG Antibody, Rabbit PAb, Antigen Affinity Purified, Immunofluorescence

Immunofluorescence staining of UNG in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-human UNG polyclonal antibody (1  $\mu$ g/ml) at 4°C overnight. Then cells were stai



sections).