

# Anti-Olig-2 antibody



Catalog Number: 101184

## Product name

Anti-Olig-2 antibody

WB: 2-5 µg/mL

IHC-P: 1-10 µg/mL

## Specificity

Human Olig-2 / Oligo2


## Validations

## Antibody description

Rabbit monoclonal to Olig-2

## Preparation

This antibody was obtained from a rabbit immunized with purified, synthetic peptide corresponding to the N-terminus of the Human Olig-2.



Olig-2 / Oligo2 Antibody, Rabbit MAb, Immunohistochemistry

## Formulation

0.2 µm filtered solution in PBS with 5% trehalose


Immunochemical staining of human olig-2 in human brain with rabbit monoclonal antibody (5 µg/mL, formalin-fixed paraffin embedded

## Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C.

Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.



sections).

Olig-2 / Oligo2 Antibody, Rabbit MAb, Immunohistochemistry

## Clonality

Monoclonal

## Ig Type

Rabbit IgG

Immunochemical staining of human olig-2 in human glioma with rabbit monoclonal antibody (5 µg/mL, formalin-fixed paraffin embedded

## Applications


WB, IHC-P

## Dilutions

# Anti-Olig-2 antibody



Catalog Number: 101184



Olig-2 / Oligo2 Antibody, Rabbit MAb, Western blot

sections).

Items	Lanes	A
Sample (whole cell lysate)		Jurkat
Sample Volume ( $\mu$ g/lane)		30
Gel		13% SDS-PAGE reducing gel
Recommended Concentration		2-5 $\mu$ g/ml
Secondary Antibody		DyLight 800-labeled Antibody To Rabbit IgG (H+L), at 1:5000 dilution.