

# Anti-CXADR / CAR antibody



Catalog Number: 104680

## Product name

Anti-CXADR / CAR antibody

## Immunogen

[Human CXADR / CAR \(His Tag\) recombinant protein](#)

## Specificity

Human CXADR / CAR

## Antibody description

Rabbit monoclonal to CXADR / CAR

## Preparation

This antibody was obtained from a rabbit immunized with purified, recombinant Human CXADR / CAR (rh CXADR / CAR; NP\_001329.1; Met1-Gly237).

## Formulation

0.2  $\mu$ m filtered solution in PBS with 5% trehalose

## Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

## Clonality

Monoclonal

## Ig Type

Rabbit IgG

## Applications


FCM, IF, ICC/IF

## Dilutions

FCM: 0.5-2  $\mu$ g/Test


ICC/IF: 10-25  $\mu$ g/mL

## Validations



CXADR / CAR Antibody, Rabbit MAb, Flow cytometric analysis

Flow cytometric analysis of Human CXADR expression on HT-29 cells. Cells were stained with purified anti-Human CXADR, then a FITC-conjugated second step antibody. The histogram were derived from gated events with the forward and side light-scatter characteristics of intact



CXADR / CAR Antibody, Rabbit MAb

Immunofluorescence staining of Human CXADR in U2OS cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with

## Anti-CXADR / CAR antibody



Catalog Number: 104680

---

rabbit anti-Human CXADR monoclonal antibody (15 µg/ml) at 4°C overnight. Then cells were stained with the Alexa Fluor® 594-conjugated

Goat Anti-rabbit IgG secondary antibody (red) and counterstained with DAPI (blue). Positive staining was localized to cells membrane.