# Product name

Anti-NBL1/DAND1 antibody

## Immunogen

Human NBL1/DAND1 (His Tag) recombinant protein

## Specificity

Human NBL1 / DAND1 / DAN

## **Antibody description**

Mouse monoclonal to NBL1/DAND1

## Preparation

This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human NBL1 / DAND1 / DAN (rh NBL1 / DAND1 / DAN; NP\_005371.1; Met1-Asp180). The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.

# Formulation

 $0.2\ \mu m$  filtered solution in PBS with 5% trehalose

#### Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

# Clonality

Monoclonal

Ig Type

Mouse IgG1


# Applications

IHC-P

## Dilutions


IHC-P: 5-20 µg/mL

## Validations



NBL1 / DAND1 / DAN Antibody, Mouse MAb, Immunohistochemistry

Immunochemical staining of human DAN in human kidney with mouse monoclonal antibody (10 µg/mL, formalin-fixed paraffin embedded sections). The left panel: tissue incubated with primary antibody; The right panel: tissue incubated with the mixture of primary antibody and antigen (recombinant protein).




NBL1 / DAND1 / DAN Antibody, Mouse MAb, Immunohistochemistry

Immunochemical staining of human DAN in human liver with mouse monoclonal antibody (10  $\mu$ g/mL, formalin-fixed paraffin embedded sections). The left panel: tissue incubated with primary antibody; The right panel: tissue





incubated with the mixture of primary antibody and antigen (recombinant protein).



DAN Antigen

NBL1 / DAND1 / DAN Antibody, Mouse MAb, Immunohistochemistry

Immunochemical staining of human DAN in human lung with mouse monoclonal antibody (10 µg/mL, formalin-fixed paraffin embedded sections). The left panel: tissue incubated with primary antibody; The right panel: tissue incubated with the mixture of primary antibody and antigen (recombinant protein).