# Anti-Cadherin-17/LI-cadherin antibody

Catalog Number: 106425



#### **Product name**

Anti-Cadherin-17/LI-cadherin antibody

#### **Immunogen**

<u>Human Cadherin-17/LI-cadherin (His Tag)</u> <u>recombinant protein</u>

# **Specificity**

Human Cadherin-17 / LI-cadherin / CDH17 **No cross-reactivity** in ELISA with Human Cad8; Human E-Cad; Human Cad5

## Antibody description

Rabbit monoclonal to Cadherin-17/LI-cadherin

# **Preparation**

This antibody was obtained from a rabbit immunized with purified, recombinant Human Cadherin-17 / LI-cadherin / CDH17 (rh Cadherin-17 / LI-cadherin / CDH17; Q12864-1; Met1-Met787).

#### **Formulation**

0.2 µm filtered solution in PBS with 5% trehalose

#### Storage

This antibody can be stored at  $2^{\circ}\text{C-8°C}$  for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C.

Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

#### **Clonality**

Monoclonal

# Ig Type

Rabbit IgG

# **Applications**

IHC-P

#### **Dilutions**

IHC-P: 1-10 μg/mL

#### **Validations**



Cadherin-17 / LI-cadherin / CDH17 Antibody, Rabbit MAb, Immunohistochemistry

Immunochemical staining of human CDH17 in human appendix with rabbit monoclonal antibody (5  $\mu$ g/mL, formalin-fixed paraffin embedded sections). Positive staining was localized to intestinal gland.



Cadherin-17 / LI-cadherin / CDH17 Antibody, Rabbit MAb, Immunohistochemistry


Immunochemical staining of human CDH17 in human colon with rabbit monoclonal antibody (5

# Anti-Cadherin-17/LI-cadherin antibody

Catalog Number: 106425



µg/mL, formalin-fixed paraffin embedded sections). Positive staining was localized to colonic



Cadherin-17 / LI-cadherin / CDH17 Antibody, Rabbit MAb, Immunohistochemistry

Immunochemical staining of human CDH17 in human rectal carcinoma with rabbit monoclonal antibody (5  $\mu$ g/mL, formalin-fixed paraffin embedded sections). Positive staining was localized to intestinal gland.