

Anti-Biotinidase / biotinase / BTD antibody



Catalog Number: 106689

Product name

Anti-Biotinidase / biotinase / BTD antibody

Immunogen

[Human Biotinidase / biotinase / BTD \(His Tag\) recombinant protein](#)

Specificity

Human Biotinidase / biotinase / BTD

Antibody description

Rabbit polyclonal to Biotinidase / biotinase / BTD

Preparation

Produced in rabbits immunized with purified, recombinant Human Biotinidase / biotinase / BTD (rh Biotinidase / biotinase / BTD; P43251; Met1-Asp543). Biotinidase / biotinase / BTD specific IgG was purified by Human Biotinidase / biotinase / BTD affinity chromatography.

Formulation

0.2 µm filtered solution in PBS

Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C.

Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

Clonality

Polyclonal

Ig Type

Rabbit IgG

Applications

ELISA, WB, IHC-P

Dilutions


WB: 5-10 µg/ml

ELISA:0.1-0.2 µg/mL

This antibody can be used at 0.1-0.2 µg/mL with the appropriate secondary reagents to detect Human Biotinidase / biotinase / BTD. The detection limit for Human Biotinidase / biotinase / BTD is < 0.039 ng/well.


IHC-P: 0.1-2 µg/ml

Validations




Lanes	A
Items	
Sample (whole cell lysate)	Jurkat
Sample Volume (µg/lane)	30
Gel	13% SDS-PAGE reducing gel
Recommended Concentration	5-10 µg/ml
Secondary Antibody	Dylight 800-labeled Antibody To Rabbit IgG (H+L), at 1:5000 dilution

Biotinidase / biotinase / BTD Antibody, Rabbit PAb, Antigen Affinity Purified, Western blot



Biotinidase / biotinase / BTD Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry



Biotinidase / biotinase / BTD Antibody, Rabbit
PAb, Antigen Affinity Purified,
Immunohistochemistry