

Anti-HDAC1 antibody



Catalog Number: 106729

Product name

Anti-HDAC1 antibody

IHC-P: 1-5 µg/mL

ICC/IF: 0.5-1.5 µg/mL


Specificity

Human HDAC1

Validations

Antibody description

Rabbit polyclonal to HDAC1



HDAC1 Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Preparation

Produced in rabbits immunized with a synthetic peptide corresponding to the center region of the Human HDAC1, and purified by antigen affinity chromatography.

Immunochemical staining of human HDAC1 in human tonsil with rabbit polyclonal antibody (2 µg/mL, formalin-fixed paraffin embedded

Formulation


0.2 µm filtered solution in PBS

Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C.

Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.



sections).

HDAC1 Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Clonality

Polyclonal

Ig Type

Rabbit IgG

Immunochemical staining of human HDAC1 in human testis with rabbit polyclonal antibody (2 µg/mL, formalin-fixed paraffin embedded

Applications

IHC-P, IF, ICC/IF

Dilutions


Anti-HDAC1 antibody



Catalog Number: 106729

HDAC1 Antibody, Rabbit PAb, Antigen Affinity Purified, Immunofluorescence

Immunofluorescence staining of HDAC1 in HeLa cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-human HDAC1 polyclonal antibody (1 $\mu\text{g}/\text{ml}$) at 4°C overnight. Then cells were



sections).