

**Product name**

Anti-SCN9A antibody

**Specificity**

Human

**Antibody description**

Rabbit polyclonal antibody to SCN9A

**Preparation**

This antigen of this antibody was synthetic peptide within human nav1.7 aa 1-50 (cytoplasmic).

**Formulation**

Liquid, 1\*PBS (pH7.4), 0.2% BSA, 50% Glycerol.  
Preservative: 0.05% Sodium Azide.

**Storage**

Store at +4°C after thawing. Aliquot store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

**Clonality**

Polyclonal

**Ig Type**

Rabbit IgG

**Applications**

ICC, IHC-P, FC

**Dilutions**

ICC: 1:50-1:200

ICC: 1:50-1:200

FC: 1:50-1:100


**Validations**

Fig1: ICC staining NaV1.7 in A549 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Fig2: ICC staining NaV1.7 in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Fig3: ICC staining NaV1.7 in SH-SY5Y cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.




Fig4: Immunohistochemical analysis of paraffin-embedded human liver cancer tissue using anti-NaV1.7 beta antibody. Counter stained with

hematoxylin.




Fig5: Immunohistochemical analysis of paraffin-embedded mouse colon tissue using anti-NaV1.7 beta antibody. Counter stained with hematoxylin.




Fig6: Flow cytometric analysis of SH-SY5Y cells with NaV1.7 antibody at 1/100 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black).