# **Anti-NCR3LG1 antibody**

Catalog Number: 175211



#### Product name

Anti-NCR3LG1 antibody

# **Specificity**

Human, Mouse

# **Antibody description**

Rabbit polyclonal antibody to NCR3LG1

# Preparation

This antigen of this antibody was klh conjugated synthetic peptide derived from human b7-h6 51-150/454

#### **Formulation**

Liquid, 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

### **Storage**

Store at -20°C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4°C.

## **Clonality**

Polyclonal

## Ig Type

IgG

## **Applications**

WB, IHC-P

#### **Dilutions**

WB:1:500-2000

IHC-P:1:400-800

#### **Validations**





Fig1: Paraformaldehyde-fixed, paraffin embedded (Human brain glioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (B7-H6) Polyclonal Antibody, Unconjugated at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023)




instructions and DAB staining.

Fig2: Sample:; Huh7 Cell (Human) Lysate at 40 ug; A549 Cell (Human) Lysate at 40 ug; Primary: Anti-B7-H6 at 1/300 dilution; Secondary: HRP conjugated Goat-Anti-rabbit IgG (bs-0295G-HRP) at 1/5000 dilution; Predicted band size: 48 kD;

# **Anti-NCR3LG1 antibody**

Catalog Number: 175211





Observed band size: 48 kD

Fig3: Sample:; Heart (Mouse) Lysate at 40 ug; Liver (Mouse) Lysate at 40 ug; Primary: Anti- B7-H6 at 1/1000 dilution; Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution; Predicted band size: 48 kD; Observed band size:




Fig4: Sample:; Cerebral cortex (Mouse) Lysate at 40 ug; Hippocampus (Mouse) Lysate at 40 ug; Primary: Anti- B7-H6 at 1/1000 dilution; Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution; Predicted band size: 48 kD; Observed band size: 50 kD