Anti-HBP1 antibody

NovoPro

Product name

Anti-HBP1 antibody

Specificity

Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Horse, Rabbit, Sheep

Antibody description

Rabbit polyclonal antibody to HBP1

Preparation

This antigen of this antibody was klh conjugated synthetic peptide derived from human hbp1 421-514/514

Formulation

Liquid, 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

Storage

Store at -20°C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4°C.

Clonality

Polyclonal

Ig Type

Rabbit IgG

Applications

WB, IHC-P

Dilutions

WB:1:500-2000

IHC-P:1:400-800

Validations

		12%Gel	
120kD			
85kD			
50kD	-	58kD	
35kD			
25kD			
20kD			

Fig1: Protein: intestine(mouse) lysates at 40ug;; Primary: Anti-HBP1 at 1:200;; Secondary: HRP conjugated Goat Anti-Rabbit IgG(bs-0295G-HRP) at 1: 3000;; ECL excitated the fluorescence;; Predicted band size : 58kD; Observed band size :




Fig2: Sample:; placenta (mouse) Lysate at 40 ug; Uterus (Mouse) Lysate at 40 ug; bone (Mouse) Lysate at 40 ug; Primary: Anti-HBP1 at 1/300 dilution; Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution; Predicted band size: 58 kD; Observed band size: 63 kD

Anti-HBP1 antibody

Catalog Number: 175489






Fig3: Paraformaldehyde-fixed, paraffin embedded (Mouse intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HBP1) Polyclonal Antibody, Unconjugated at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.