

Anti-IL1rn antibody

Catalog Number: 175517

Product name

Anti-IL1rn antibody

Specificity

Human, Mouse, Rat, Dog, Pig, Cow, Sheep

Antibody description

Rabbit polyclonal antibody to IL1rn

Preparation

This antigen of this antibody was klh conjugated synthetic peptide derived from human il-1ra 26-120/178

Formulation

Liquid, 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

Storage

Store at -20°C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4°C.

Clonality

Polyclonal

Ig Type

IgG

Applications

WB, IHC-P

Dilutions

WB:1:500-2000

IHC-P:1:400-800

Validations





Fig1: Tissue/cell: Human oral squamous cell carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;; Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;; Incubation: Anti-IL-1RA Polyclonal Antibody, Unconjugated 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023)



and DAB(C-0010) staining

Fig2: Sample: Lane 1:Plasma(Mouse) Tissue Lysate at 30 ug; Lane 2:Hemocyte(Mouse) Tissue Lysate at 30 ug; Primary: Anti-IL-1RA at 1/300 dilution; Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution; Predicted band size: 18 kD; Observed band size: 23 kD

Catalog Number: 175517





Fig3: Sample: Spleen(Mouse) Tissue Lysate at 30 ug; Primary: Anti-IL-1RA at 1/300 dilution; Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution; Predicted band size: 18 kD; Observed band size: 23 kD

Fig4: Paraformaldehyde-fixed, paraffin embedded (mouse spinal tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (IL-1RA) Polyclonal Antibody, Unconjugated at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.