

Human VEGFR2/Flk-1/CD309/KDR (Fc Tag) recombinant protein



Catalog Number: 501107

General Information

Protein Construction

A DNA sequence encoding the extracellular domain of human VEGFR2 (NP_002244.1) (Met 1-Glu 764) was fused with the Fc region of human IgG1 at the C-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

1. Measured by its binding ability in a functional ELISA.
Immobilized recombinant human VEGFA at 1.25 µg/ml (100 µl/well) can bind VEGFR2 with a linear range of 1.25-40.0 ng/ml.
2. Measured by its ability to inhibit the VEGF-dependent proliferation of human umbilical vein endothelial cells (HUVEC). The ED₅₀ for this effect is typically 10-40 ng/mL in the presence of 10 ng/mL recombinant human VEGF165.

Purity

> 97 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Ala 20

Molecular Mass

The recombinant human VEGFR2/Fc is a disulfide-linked homodimeric protein. The reduced monomer consists of 979 amino acids and predicts a molecular mass of 109 kDa. The apparent molecular mass of rh VEGFR2/Fc monomer is approximately 150-160 in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.
2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human VEGFR2 / Flk-1 / CD309 / KDR Protein (Fc Tag) SDS-PAGE