

Anti-SPARCL1 / SPARC-like 1 antibody



Catalog Number: 106267

Product name

Anti-SPARCL1 / SPARC-like 1 antibody

Specificity

Human SPARCL1 / SPARC-like 1

Antibody description

Mouse monoclonal to SPARCL1 / SPARC-like 1

Preparation

This antibody was produced from a hybridoma resulting from the fusion of a mouse myeloma with B cells obtained from a mouse immunized with purified, recombinant Human SPARCL1 / SPARC-like 1 (rh SPARCL1 / SPARC-like 1; NP_004675.3; Met1-Phe664). The IgG fraction of the cell culture supernatant was purified by Protein A affinity chromatography.

Formulation

0.2 μ m filtered solution in PBS with 5% trehalose

Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C.

Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

Clonality

Monoclonal

Ig Type

Mouse IgG2b

Applications


FCM, IF, ICC/IF

Dilutions

FCM: 0.5-2 μ g/Test


ICC/IF: 10-25 μ g/mL

Validations



SPARCL1 / SPARC-like 1 Antibody, Mouse MAb, Flow cytometric analysis

Flow cytometric analysis of Human SPARCL1 expression on HL60 cells. The cells were treated according to manufacturer's manual (BD Pharmingen™ Cat. No. 554714), stained with purified anti-Human SPARCL1, then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.



SPARCL1 / SPARC-like 1 Antibody, Mouse MAb, Immunofluorescence

Immunofluorescence staining of Human SPARCL1 in A431 or HeLa cells. Cells were fixed with 4% PFA, permeabilized with 1% Triton X-100 in PBS, blocked with 10% serum, and incubated with

Anti-SPARCL1 / SPARC-like 1 antibody



Catalog Number: 106267

mouse anti-Human SPARCL1 monoclonal antibody (15 µg/ml). Then cells were stained with the Alexa Fluor® 488-conjugated Goat Anti-mouse IgG secondary antibody (left panel, captured by laser

confocal scanning microscope; right panel, captured by fluorescence microscope), counterstained with DAPI (blue). Positive staining was localized to cytoplasm.