

Human CD97 (Fc Tag) recombinant protein



Catalog Number: 504197

General Information

Gene Name Synonym

Leukocyte antigen CD97; CD97 antigen subunit alpha; CD97 antigen subunit beta

Protein Construction

A DNA sequence encoding the first 398 amino acids (Met 1-Gln 398) of human CD97 isoform 2 (NP_001775.2) extracellular domain was fused with the Fc region of human IgG1 at the C-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Measured by its binding ability in a functional ELISA. Immobilized human CD55 at 2 µg/ml (100 µl/well) can bind human CD97 with a linear range of 1.28-32 ng/ml.

Purity

> 90 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Gln 21

Molecular Mass

The recombinant human CD97/Fc chimera is a disulfide-linked homodimeric protein. The reduced monomer consists of 619 amino acids and predicts a molecular mass of 68.2 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rh CD97/Fc monomer is approximately 100-110 kDa due to glycosylation.

Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.
2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human CD97 Protein (Fc Tag) SDS-PAGE