Catalog Number: 500204



General Information

Gene Name Synonym

Macrophage inhibitory cytokine 1; NSAIDactivated gene 1 protein; NSAID-regulated gene 1 protein; Placental TGF-beta; Placental bone morphogenetic protein; Prostate differentiation factor

Protein Construction

A DNA sequence encoding the mature form of human GDF15 (NP_004855.2) (Ala 197-Ile 308) was expressed with the fused Fc region of human IgG1 at the N-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Purity

> 95 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70 $^\circ \rm C$

Predicted N terminal

Glu 20

Molecular Mass

The recombinant human Fc/GDF15 is a disulfidelinked homodimeric protein. The reduced monomer consists of 370 amino acids and has a predicted molecular mass of 40.6 kDa. rhGDF15/Fc monomer migrates as an approximately 44 kDa protein in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile 100mM Glycine, 10mM NaCl, 50mM Tris, pH 7.5 1. 5 % trehalose and mannitol are added as protectants before lyophilization. 2. Please contact us for any concerns or special requirements.

Usage Guide

Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE

KDa	М	
116		
66.2		
45.0		-
35.0	-	
25.0	-	
18.4	-	
14.4	-	

Human GDF-15 Protein (Fc Tag) SDS-PAGE