Human IL-10 (His Tag) recombinant protein

Catalog Number: 500745



General Information

Gene Name Synonym

Cytokine synthesis inhibitory factor

Protein Construction

A DNA sequence encoding the human IL10 (NP_000563.1) (Ser 19-Asn 178) was expressed, with a N-terminal polyhistidine tag.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

- 1. Immobilized human IL10-His at 10 μ g/ml (100 μ l/well) can bind Cynomolgus IL10RA-Fc (Cat:504750), The EC₅₀ of Cynomolgus IL10RA-Fc (Cat:504750) is 0.31-0.71 μ g/ml.
- 2. Measured in a cell proliferation assay using MC/9-2 mouse mast cells. The ED_{50} for this effect is typically 5-20 ng/mL.

Purity

> 85 % as determined by SDS-PAGE

Endotoxin

 $< 1.0 \; \text{EU}$ per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

His

Molecular Mass

The recombinant human IL10 comprises 176 amino acids and has a predicted molecular mass of 20.9 kDa as estimated in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.4

- 1.5% trehalose and mannitol are added as protectants before lyophilization.
- 2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human IL10 / Interleukin-10 Protein (His Tag) SDS-PAGE