

Human DOPA Decarboxylase/DDC (His Tag) recombinant protein



Catalog Number: 500922

General Information

Gene Name Synonym

DOPA decarboxylase

Protein Construction

A DNA sequence encoding the full length of human DDC (NP_000781.1) (Met 1-Glu 480) was expressed with a polyhistidine tag at the C-terminus.

Organism

Human

Expression Host

Baculovirus-Insect Cells

QC Testing

Activity

Measured by its ability to convert the substrate 3, 4-dihydroxy L-phenylalanine (L-Dopa) to 3, 4-dihydroxyphenylethylamine (dopamine). The dopamine product is measured by its absorbance at 340 nm after derivatization with trinitrobenzene sulfonic acid. The specific activity is >1500 pmoles/min/μg.

Purity

> 90 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Met

Molecular Mass

The recombinant human DDC consists of 490 amino acids and predicts a molecular mass of 55 kDa. It migrates as an approximately 48 kDa protein in SDS-PAGE under reducing conditions.

Formulation

Supplied as sterile 50mM Tris, 100mM NaCl, pH 8, 10% glycerol

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human DOPA Decarboxylase / DDC Protein (His Tag) SDS-PAGE