

General Information

Gene Name Synonym

Interferon alpha binding protein; Type I interferon receptor 2

Protein Construction

A DNA sequence encoding the extracellular domain of human IFNAR2 isoform a (NP_997468.1) (Met 1-Lys 243) was fused with the Fc region of human IgG1 at the C-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Measured by its ability to inhibit rh IFN β mediated protection of WISH Human amnion cells infected with vesicular stomatitis virus (VSV) to viral lysis.

The EC₅₀ for this effect is typically 0.01-0.05 μ g/mL.

Purity

> 90 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μ g of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Ile 27

Molecular Mass

The recombinant human IFNAR2/Fc is a disulfide-linked homodimeric protein. The reduced monomer consists of 458 amino acids and predicts a molecular mass of 51.8 kDa. As a result of glycosylation, the apparent molecular mass of rh IFNAR2/Fc monomer is approximately 65-75 kDa in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.
2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human IFNAR2 / IFNABR Protein (Fc Tag) SDS-

Human IFNAR2 (Fc Tag) recombinant protein



Catalog Number: 501243

PAGE