

General Information

Gene Name Synonym

Proto-oncogene c-Mer; Receptor tyrosine kinase
MerTK

Protein Construction

A DNA sequence encoding the mouse MERTK (Q60805) extracellular domain (Met 1-Phe 498) was fused with the Fc region of human IgG1 at the C-terminus.

Organism

Mouse

Expression Host

Human Cells

QC Testing

Purity

> 90 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Gly 19

Molecular Mass

The secreted recombinant mouse MERTK/Fc

chimera is a disulfide-linked homodimer. The reduced monomer consists of 721 amino acids and has a predicted molecular mass of 79.3 kDa. As a result of glycosylation, rm MERTK/Fc monomer migrates as an approximately 130 kDa band in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C . It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Mouse MERTK / Mer Protein (Fc Tag) SDS-PAGE