Catalog Number: 501764



General Information

Gene Name Synonym

Fallotein; Spinal cord-derived growth factor; VEGF-E; Platelet-derived growth factor C, latent form; Platelet-derived growth factor C, receptorbinding form

Protein Construction

A DNA sequence encoding the receptor-binding form of mouse PDGFC (Q8CI19) (Val 235-Gly 345) was fused with the Fc region of human IgG1 at the N-terminus.

Organism

Mouse

Expression Host

Human Cells

QC Testing

Activity

1. Measured by its binding ability in a functional ELISA.

2. Immobilized mouse PDGF-C (Cat:501764) at 10 μ g/mL (100 μ l/well) can bind human PDGFRA (Cat:503138), The EC₅₀ of human PDGFRA (Cat:503138) is 0.71 μ g/mL.

3. Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast cells. The ED_{50} for this effect is typically 0.5-2 µg/mL.

Purity

> 80 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70 $^{\circ}\mathrm{C}$

Predicted N terminal

Glu

Molecular Mass

The recombinant mouse PDGFC/Fc is a disulfidelinked homodimeric protein. The reduced monomer consists of 372 amino acids and has a predicted molecular mass of 40.9 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rmPDGFC/Fc monomer is approximately 46 kDa.

Formulation

Lyophilized from sterile PBS, pH 7.4 1. 5 % trehalose and mannitol are added as protectants before lyophilization. 2. Please contact us for any concerns or special requirements.

Usage Guide

Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.


Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Catalog Number: 501764



Mouse PDGF-C / SCDGF Protein (Fc Tag) SDS-PAGE