Human CXADR / CAR (His Tag) recombinant protein

Catalog Number: 502086

General Information

Gene Name Synonym

CVB3-binding protein; Coxsackievirus Badenovirus receptor; HCVADR

Protein Construction

A DNA sequence encoding the human CXADR (NP_001329.1) extracellular domain (Met 1-Gly 237) was fused with a polyhistidine tag at the C-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Measured by the ability of the immobilized protein to support the adhesion of mouse neutrophils. When 5 x 10E4 cells/well are added to CXADR coated plates (4 μ g/ml and 100 μ l/well), approximately 20%-40% will adhere specifically after 60 minutes at 37 °C.

Purity

> 92 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70 $^{\circ}\mathrm{C}$

Predicted N terminal

Leu 20

Molecular Mass

The secreted recombinant human CXADR consists of 229 amino acids and has a predicted molecular mass of 25.6 kDa. As a result of glycosylation, the apparent molecular mass of rh CXADR is approximately 35 kDa in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.41. 5 % trehalose and mannitol are added as protectants before lyophilization.2. Please contact us for any concerns or special requirements.

Usage Guide

Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE

KDa	М	
116	-	
66.2	-	
45.0	-	
35.0		,
25.0	-	
18.4	-	
14.4	-	

Human CXADR / CAR Protein (His Tag) SDS-PAGE

