Catalog Number: 502476



General Information

Gene Name Synonym

TGF-beta type II receptor; Transforming growth factor-beta receptor type II

Protein Construction

A DNA sequence encoding the rhesus TGFBR2 (NP_001248080.1) (Met1-Arg159) was expressed with the Fc region of human IgG1 at the C-terminus.

Organism

Rhesus

Expression Host

Human Cells

QC Testing

Activity

1. Measured by its binding ability in a functional ELISA.

2. Immobilized Rhesus TGFB1-His (Cat:504900) at 10 $\mu\text{g/mL}$ (100 $\mu\text{L/well})$ can bind Rhesus TGFBR2-Fc.

The EC_{50} of Rhesus TGFBR2-Fc is 28-56 ng/mL.

Purity

> 95 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70 $^{\circ}\mathrm{C}$

Predicted N terminal Ile 24

Molecular Mass

The recombinant rhesus TGFBR2 comprises 377 amino acids and has a calculated molecular mass of 42.4 KDa.

Formulation

Lyophilized from sterile PBS, pH 7.41. 5 % trehalose and mannitol are added as protectants before lyophilization.2. Please contact us for any concerns or special requirements.

Usage Guide

Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE

KDa	M
116	-
66.2	- 65
45.0	-
35.0	-
25.0	-
18.4	-
14.4	-

Cynomolgus TGFBR2 Protein (Fc Tag) SDS-PAGE