

Human ICOS Ligand/B7-H2 (Fc Tag) recombinant protein



Catalog Number: 502558

General Information

Gene Name Synonym

B7 homolog 2; B7-like protein G150; B7-related protein 1

Protein Construction

A DNA sequence encoding the human ICOSLG (O75144-1) (Met1-Ser258) was expressed with the Fc region of human IgG1 at the C-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Purity

(96.1±2.0) % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Asp 19

Molecular Mass

The recombinant human ICOSLG/Fc is a disulfide-linked homodimer. The reduced monomer

comprises 481 amino acids and has a predicted molecular mass of 53.7 kDa. The apparent molecular mass of the protein is approximately 66-76 and 33 kDa in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.4.

1. 5 % trehalose and mannitol are added as protectants before lyophilization.
2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human ICOS Ligand / B7-H2 / ICOSLG Protein (Fc Tag) SDS-PAGE