

Human Cystatin 7 / CST7 (Fc Tag) recombinant protein



Catalog Number: 503019

General Information

Gene Name Synonym

Cystatin-7; Cystatin-like metastasis-associated protein; Leukocystatin

Protein Construction

A DNA sequence encoding the human Cystatin F precursor (NP_003641.3) (Met 1-His 145) was expressed with the fused Fc region of human IgG1 at the C-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Measured by its ability to inhibit active Cathepsin L. Cleavage of a fluorogenic peptide substrate Z-LR-AMC (R&D Systems, Catalog#ES008). The IC₅₀ value is <5 nM.

Purity

> 97 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Gly 20

Molecular Mass

The recombinant human Cystatin-F/Fc is a disulfide-linked homodimeric protein. The reduced monomer consists of 364 amino acids and predicts a molecular mass of 41.3 kDa. As a result of glycosylation, the rh Cystatin-F/Fc monomer migrates as an approximately 50 kDa band in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile 100mM Glycine, 10mM NaCl, 50mM Tris, pH 7.5

1. 5 % trehalose and mannitol are added as protectants before lyophilization.
2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human Cystatin 7 / CST7 Protein (Fc Tag) SDS-PAGE