Human IL18BP / IL18BPa (His & Fc Tag) recombinant protein

Catalog Number: 503436

General Information

Protein Construction

A DNA sequence encoding the N-terminal domain (Met 1-Gly 192) of human IL 18BPa (NP_001034748.1) was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Measured by its ability to bind Cynomolgus IL18 in functional ELISA.

Purity

> 95 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70 $^{\circ}\mathrm{C}$

Predicted N terminal

Thr 29

Molecular Mass

The recombinant human IL18BPa/Fc is a disulfidelinked homodimeric protein after removal of the



signal peptide. The reduced monomer consists of 411 amino acids and has a predicted molecular mass of 44.5 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rh IL18BPa/Fc monomer is approximately 70-75 kDa due to glycosylation.

Formulation

Lyophilized from sterile 100mM Glycine, 10mM NaCl, 50mM Tris, pH 7.5 1. 5 % trehalose and mannitol are added as protectants before lyophilization. 2. Please contact us for any concerns or special requirements.

Usage Guide

Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE

KDa	М	
116		
66.2	-	μ
45.0	-	
35.0	-	
25.0	-	
18.4	_	
14.4	-	

Human IL18BPa Protein (His & Fc Tag) SDS-PAGE