Human Thrombopoietin/THPO (His Tag) recombinant protein

Catalog Number: 503688

General Information

Gene Name Synonym

C-mpl ligand; Megakaryocyte colony-stimulating factor; Megakaryocyte growth and development factor; Myeloproliferative leukemia virus oncogene ligand

Protein Construction

A DNA sequence encoding the mature form of human THPO (NP_000451.1) (Ser 22-Gly 353) was fused with a polyhistidine tag at the N-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Measured in a cell proliferation assay using MO7e human megakaryocytic leukemic cells. The ED_{50} for this effect is typically 3-15 ng/ml.

Purity

> 95 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70 $^{\circ}\mathrm{C}$

Predicted N terminal

Ser 22 **Molecular Mass**

The recombinant human THPO consists of 348 amino acids with the predicted molecular mass of 37.6 kDa. As a result of glycosylation, rhTHPO migrates as an approximately 80 kDa band in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.41. 5 % trehalose and mannitol are added as protectants before lyophilization.2. Please contact us for any concerns or special requirements.

Usage Guide

Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE

KDa	М	
116		
66.2	-	
45.0	-	
35.0	-	
25.0	-	
18.4	-	
14.4	-	

Human Thrombopoietin / THPO / TPO Protein (His Tag) SDS-PAGE

