Human CTLA-4/CD152 (His Tag) recombinant protein

Catalog Number: 503821

NovoPro

General Information

Gene Name Synonym

Cytotoxic T-lymphocyte-associated antigen 4

Protein Construction

A DNA sequence encoding the human CTLA4 (NP_005205.2) extracellular domain (Met 1-Phe 162) was fused with the a polyhistidine tag at the C-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

1. Measured by its binding ability in a functional ELISA.

2. Immobilized human CTLA4 (cat: 503821) at 10 μ g/mL (100 μ l/well) can bind human B7-1/Fc (cat: 503507), The EC₅₀ of human B7-1/Fc (cat: 503507 is 14 ng/mL.

3. Immobilized human CTLA4 (cat: 503821) at 10 μ g/mL (100 μ l/well) can bind human B7-2/Fch (cat: 504521), The EC₅₀ of human B7-2/Fch (cat: 504521) is 53 ng/mL.

Purity

> 90 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from

date of receipt at -70°C Predicted N terminal

Ala 37

Molecular Mass

The recombinant human CTLA4 consists of 137 amino acids and has a predicted molecular mass of 15 kDa.

Formulation

Lyophilized from sterile PBS, pH 7.41. 5 % trehalose and mannitol are added as protectants before lyophilization.2. Please contact us for any concerns or special requirements.

Usage Guide

Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE

KDa	M
116	
66.2	-
45.0	-
35.0	-
25.0	- 665
18.4	-
14.4	-

Human CTLA4 / CD152 Protein (His Tag) SDS-PAGE