Mouse VCAM-1/CD106 (His & Fc Tag) recombinant protein

Catalog Number: 503920



General Information

Protein Construction

A DNA sequence encoding the extracellular domain (Met 1-Glu 698) of mouse VCAM1 (NP_035823.3) precursor was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.

Organism

Mouse

Expression Host

Human Cells

QC Testing

Activity

Measured by the ability of the immobilized protein to support the adhesion of U937 human histiocytic lymphoma cells.

When cells are added to VCAM1 coated plates (10 μ g/ml, 100 μ g/well) approximately >70% cells will adhere specifically.

Purity

> 90 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Phe 25

Molecular Mass

The recombinant mouse VCAM1/Fc is a disulfide-linked homodimer after removal of the signal peptide. The reduced monomer consists of 922 amino acids and has a predicted molecular mass of 102 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rmVCAM1/Fc monomer is approximately 110-120 kDa due to glycosylation.

Formulation

Lyophilized from sterile PBS, pH 7.4

- 1.5% trehalose and mannitol are added as protectants before lyophilization.
- 2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Mouse VCAM1 / CD106 Protein (His & Fc Tag) SDS-PAGE