Mouse CD132 / IL2RG (His & Fc Tag) recombinant protein

Catalog Number: 504271

General Information

Gene Name Synonym

Interleukin-2 receptor subunit gamma; gammaC; p64

Protein Construction

A DNA sequence encoding the extracellular domain (Met 1-Ala 263) of mouse IL2RG (NP_038591.1) precursor was fused with the Cterminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.

Organism

Mouse

Expression Host

Human Cells

QC Testing

Activity

Measured by its ability to bind biotinylated recombinant rat IL2 in a functional ELISA.

Purity

> 92 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70 $^{\circ}\mathrm{C}$

Predicted N terminal

Trp 23

Molecular Mass

The recombinant mouse IL2RG/Fc is a disulfidelinked homodimer after removal of the signal peptide. The reduced monomer consists of 489 amino acids and has a predicted molecular mass of 56.3 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rm IL2RG/Fc monomer is approximately 75-85 kDa due to glycosylation.

Formulation

Lyophilized from sterile PBS, pH 7.41. 5 % trehalose and mannitol are added as protectants before lyophilization.2. Please contact us for any concerns or special requirements.

Usage Guide

Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE

| KDa | М | |
|------|---|---|
| 116 | | - |
| 66.2 | - | - |
| 45.0 | - | |
| 35.0 | | |
| 25.0 | - | |
| 18.4 | - | 1 |
| 14.4 | _ | 1 |

Mouse IL2RG Protein (His & Fc Tag) SDS-PAGE

