

## General Information

### Gene Name Synonym

Activating signal cointegrator 2; ASC-2; Amplified in breast cancer protein 3; Cancer-amplified transcriptional coactivator ASC-2; Nuclear receptor coactivator RAP250; NRC RAP250; Nuclear receptor-activating protein, 250 kDa; Peroxisome proliferator-activated receptor-interacting protein; PPAR-interacting protein; PRIP; Thyroid hormone receptor-binding protein

### Protein Construction

A DNA sequence encoding the human NCOA6 (XP\_054179221.1) 1739-2084 aa was fused with the polyhistidine tag

### Organism

Human

### Expression Host

E. coli

## QC Testing

### Activity

Not tested.

### Endotoxin

Please contact the lab for more information.

### Stability

Store for up to 12 months at -20°C to -80°C as lyophilized powder.

### Formulation

Protein lyophilized in sterile PBS (58 mM Na<sub>2</sub>HPO<sub>4</sub>, 17 mM NaH<sub>2</sub>PO<sub>4</sub>, 68 mM NaCl, 300 mM Imidazole, pH 8.0). Trehalose (5-8%) and mannitol (5-8%) protectants were added before lyophilization.

## Usage Guide


### Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

### Reconstitution

Reconstitute at 0.25 µg/µl in sterile water for short-term storage. Reconstitution with 50% glycerol solution is recommended for longer term storage (see Stability and Storage for more details). If a different concentration is needed for your purposes please adjust the reconstitution volume as required (please note: the ion concentration of the final solution will vary according to the volume used). Note: Centrifuge vial before opening. When reconstituting, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution.

### SDS-PAGE



Recombinant human NCOA6 SDS-PAGE