

Mouse Icam2/Icam-2 (Fc tag) recombinant protein



Catalog Number: 519419

General Information

Gene Name Synonym

Intercellular adhesion molecule 2, Icam2, ICAM-2, Lymphocyte function-associated AG-1 counter-receptor, CD102, Icam-2, CD102 antigen,

Protein Construction

A DNA sequence encoding the Mouse Icam2/Icam-2 Lys23-Gln222 is expressed with a Fc tag at the C-terminus.

Organism

Mouse

Expression Host

Human Cells

QC Testing

Purity

>95% as determined by SDS-PAGE.

Endotoxin

<1.0 EU per μg protein as determined by the LAL method.

Stability

Store for up to 12 months at -20°C to -80°C as lyophilized powder. Aliquots of reconstituted samples are stable at $< -20^{\circ}\text{C}$ for 3 months.

Formulation

Lyophilized from a $0.2\ \mu\text{m}$ filtered solution of 20mM Tris, 150mM NaCl, pH8.0.

Usage Guide


Storage

Lyophilized protein should be stored at $< -20^{\circ}\text{C}$, though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at $4-7^{\circ}\text{C}$ for 2-7 days.

Reconstitution

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than $100\ \mu\text{g}/\text{ml}$. Dissolve the lyophilized protein in ddH₂O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE



Mouse Icam2/Icam-2 (Fc tag) recombinant protein