

Anti-Hnf1a antibody



Catalog Number: 175500

Product name

Anti-Hnf1a antibody

IHC-P:1:400-800

FC:3ug/test

Specificity

Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Zebrafish, Sheep

Validations

Antibody description

Rabbit polyclonal antibody to Hnf1a

Preparation

This antigen of this antibody was klh conjugated synthetic peptide derived from human hnf1 201-350/628

Formulation

Liquid, 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

Storage

Store at -20°C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4°C.

Clonality

Polyclonal

Ig Type

Rabbit IgG

Applications

WB, IHC-P, FC

Dilutions

WB:1:500-2000



Fig1: Paraformaldehyde-fixed, paraffin embedded (Mouse liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HNF1A) Polyclonal Antibody, Unconjugated at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.






Fig2: Paraformaldehyde-fixed, paraffin embedded (Rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (HNF1A) Polyclonal Antibody, Unconjugated at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023)

Anti-Hnf1a antibody

Catalog Number: 175500



instructions and DAB staining.

Fig3: Sample: HepG2 Cell (Human) Lysate at 30 μ g; Primary: Anti-HNF1 (Bs-1405R) at 1/300 dilution; Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution; Predicted band size: 67 kD; Observed band size: 67 kD

Fig4: Blank control: A549; Primary Antibody (green line): Rabbit Anti-HNF1A antibody ; Dilution: 1 μ g / 10^6 cells; Isotype Control Antibody (orange line): Rabbit IgG ; Secondary Antibody : Goat anti-rabbit IgG-PE; Dilution: 3 μ g / test; Protocol; The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.