

# Mouse ICOS Ligand/B7-H2 (His & Fc Tag) recombinant protein



Catalog Number: 504080

## General Information

### Protein Construction

A DNA sequence encoding the mouse B7-H2 (NP\_056605.1) extracellular domain (Met 1-Lys 279) was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.

### Organism

Mouse

### Expression Host

Human Cells

## QC Testing

### Activity

Measured by its binding ability in a functional ELISA. Immobilized human ICOS at 1 µg/ml (100 µl/well) can bind biotinylated mouse B7-H2 Fc chimera with a linear range of 0.125-1.0 µg/ml.

### Purity

> 95 % as determined by SDS-PAGE

### Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

### Stability

Samples are stable for up to twelve months from date of receipt at -70°C

### Predicted N terminal

Glu 47

### Molecular Mass

The secreted recombinant mouse B7-H2/Fc is a disulfide-linked homodimer after removal of the signal peptide. The reduced monomer comprises 481 amino acids with a predicted molecular mass of 54.3 kDa. As a result of glycosylation, it migrates as an approximately 75-85 kDa band in SDS-PAGE under reducing conditions.

### Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

## Usage Guide


### Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

### Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

### SDS-PAGE



Mouse ICOS Ligand / B7-H2 / ICOSLG Protein (His & Fc Tag) SDS-PAGE