

Human CD32a / Fc gamma RIIA / FCGR2A (167 Arg, His Tag) recombinant protein



Catalog Number: 503978

General Information

Gene Name Synonym

CDw32; Fc-gamma RII-a

Protein Construction

A DNA sequence encoding the extracellular domain (Met 1-Ile 218) of human CD32a (AAA35827.1) precursor was expressed, with a C-terminal polyhistidine tag. It is identical to FCGR2A131H/R in the reference.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Measured by its binding ability in a functional ELISA. Immobilized recombinant human CD32a at 10 µg/ml (100 µl/well) can bind human IgG2 with a linear range of 12.5-200 ng/ml.

Purity

> 97 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Gln 34

Molecular Mass

The recombinant human CD32a consists of 196 amino acids after removal of the signal peptide and predicts a molecular mass of 22 kDa. It migrates as an approximately 30 kDa protein due to glycosylation by SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.5

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human CD32a / FCGR2A Protein (167 Arg, His Tag) SDS-PAGE