

General Information

Gene Name Synonym

Activation B7-2 antigen; Early T-cell costimulatory molecule 1

Protein Construction

A DNA sequence encoding the extracellular domain (Met 1-Glu 245) of mouse CD86 (NP_062261.3) precursor was fused with a polyhistidine tag at the C-terminus.

Organism

Mouse

Expression Host

Human Cells

QC Testing

Activity

Measured by its binding ability in a functional ELISA. Immobilized mouse CD86 at 20 µg/ml (100 µl/well) can bind mouse CD28 with a linear ranger of 6.4-800 ng/ml.

Purity

> 92 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Val 24

Molecular Mass

The recombinant mouse CD86 consists of 233 amino acids after removal of the signal peptide and has a predicted molecular mass of 26.8 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rm CD86 is approximately 40-55 kDa due to glycosylation.

Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Mouse CD86 / B7-2 Protein (His Tag) SDS-PAGE