

General Information

Gene Name Synonym

Proto-oncogene c-Mer; Receptor tyrosine kinase MerTK

Protein Construction

A DNA sequence encoding the human MERTK (Q12866) protein kinase domain (Glu 578-Tyr 872) was fused with the N-terminal polyhistidine-tagged GST tag at the N-terminus.

Organism

Human

Expression Host

Baculovirus-Insect Cells

QC Testing

Activity

Kinase activity untested

Purity

> 92 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Met

Molecular Mass

The recombinant human MERTK (aa578-872)/GST chimera consists of 532 amino acids and has a calculated molecular mass of 62 kDa. It migrates as an approximately 50 kDa band in SDS-PAGE under reducing conditions.

Formulation

Supplied as sterile 50mM Tris, 100mM NaCl, pH 7.4, 20% gly, 0.3mM DTT

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C . It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human MERTK / Mer Protein GST Tag SDS-PAGE