# **Anti-BST1 antibody**

Catalog Number: 175221



#### **Product name**

Anti-BST1 antibody

# **Specificity**

Human, Mouse, Rat, Dog, Pig, Cow, Rabbit

## **Antibody description**

Rabbit polyclonal antibody to BST1

# **Preparation**

This antigen of this antibody was klh conjugated synthetic peptide derived from human bst1/cd157 51-150/318

#### **Formulation**

Liquid, 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

#### Storage

Store at -20°C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4°C.

# **Clonality**

Polyclonal

## Ig Type

Rabbit IgG

### **Applications**

WB, FC

#### **Dilutions**

WB:1:500-2000

FC:1µg/Test

#### **Validations**




Fig1: Blank control: U937(blue).; Primary Antibody: Rabbit Anti-BST1 antibody, Dilution: 1μg in 100 μL 1X PBS containing 0.5% BSA;; Isotype Control Antibody: Rabbit IgG (orange) ,used under the same conditions.; Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA.; Protocol; The cells were fixed with 2% paraformaldehyde (10 min). Primary antibody ( 1µg/1x10^6 cells) were incubated for 30 min on the ice, followed by 1 X PBS containing 0.5% BSA + 10% goat serum (15 min) to block non-specific protein-protein interactions. Then the Goat Antirabbit IgG/PE antibody was added into the blocking buffer mentioned above to react with the primary antibody at 1/200 dilution for 30 min on ice. Acquisition of 20,000 events was performed.




Fig2: Sample:; Raw264.7(Mouse) Cell Lysate at 30 ug; Primary: Anti-BST1 at 1/500 dilution; Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution; Predicted band size: 33 kD; Observed band size: 33 kD