

Anti-ARIH2/TRIAD1 antibody



Catalog Number: 101131

Product name

Anti-ARIH2/TRIAD1 antibody

WB: 5-10 µg/ml

Specificity

Human ARIH2

IHC-P: 0.5-3 µg/mL

ICC/IF: 0.5-1.5µg/mL


Antibody description

Rabbit polyclonal to ARIH2/TRIAD1

Validations

Preparation

Produced in rabbits immunized with a synthetic peptide corresponding to the center region of the Human ARIH2, and purified by antigen affinity chromatography.



Items	Lanes	A	B	C
Sample (whole cell lysate)		Jurkat	K562	Hela
Sample Volume (µg/lane)		30	30	30
Gel	13% SDS-PAGE reducing gel			
Recommended Concentration	5-10 µg/ml			
Secondary Antibody	Dylight 800-labeled Antibody to Rabbit IgG (H+L), at 1:5000 dilution.			
Explanation	Developed using Odyssey imaging system. Predicted band size : 57 kDa Observed band size : 57 kDa Additional bands at : 25 kDa (We are unsure as to the identity of these extra bands.)			

Formulation

0.2 µm filtered solution in PBS


ARIH2 Antibody, Rabbit PAb, Antigen Affinity Purified, Western blot

Storage

This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C.

Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.



ARIH2 Antibody, Rabbit PAb, Antigen Affinity Purified, Immunohistochemistry

Clonality

Polyclonal

Immunochemical staining of human ARIH2 in human trachea with rabbit polyclonal antibody (2 µg/mL, formalin-fixed paraffin embedded

Ig Type

Rabbit IgG

Applications


WB, IHC-P, IF, ICC/IF

Dilutions

Anti-ARIH2/TRIAD1 antibody



Catalog Number: 101131



ARIH2 Antibody, Rabbit PAb, Antigen Affinity Purified, Immunofluorescence

Immunofluorescence staining of XRCC6 in HeLa cells. Cells were fixed with 4% PFA, blocked with 10% serum, and incubated with rabbit anti-human XRCC6 polyclonal antibody (1 $\mu\text{g/ml}$) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to nucleus.

sections).