

General Information

Protein Construction

A DNA sequence encoding the mouse MET (NP_032617.2) extracellular domain (Met 1-Asn 929) was fused with the Fc region of human IgG1 at the C-terminus.

Organism

Mouse

Expression Host

Human Cells

QC Testing

Activity

Measured by its binding ability in a functional ELISA. Immobilized human HGF at 2 µg/ml (100 µl/well) can bind mouse HGFR with a linear ranger of 1.28-32ng/ml.

Purity

> 92 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Glu 25 & Ser 307

Molecular Mass

The recombinant mouse Met/Fc chimera is a disulfide-linked homodimer of the Met which is a

heterodimer composed of the proteolytically cleaved α and β subunits. Each α and β together with the C-terminal Fc tag consists of 1146 amino acids and has a predicted molecular mass of 128 (α =32 + Fc tagged β =96) kDa. The apparent molecular mass of the rm MET/Fc heterodimer thus is approximately 43 kDa and 115-120 kDa respectively in SDS-PAGE under reducing conditions due to glycosylation.

Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Mouse c-MET / HGFR Protein (Fc Tag) SDS-PAGE