

Human Nogo Receptor (His & Fc Tag) recombinant protein



Catalog Number: 503259

General Information

Gene Name Synonym

Nogo receptor; Nogo-66 receptor

Protein Construction

A DNA sequence encoding the mature form of human RTN4R (NP_075380.1) (Met 1-Ser 447) was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Measured by its ability to bind recombinant human RTN4 in a functional ELISA.

Purity

> 90 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Cys 27

Molecular Mass

The recombinant human RTN4R/Fc is a disulfide-linked homodimer. The reduced monomer consists of 669 amino acids and has a predicted molecular mass of 73 kDa. As a result of glycosylation, the apparent molecular mass of rh RTN4R/Fc monomer is approximately 100 kDa in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C . It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human Nogo Receptor / NOGOR / RTN4R Protein (His & Fc Tag) SDS-PAGE