

General Information

Gene Name Synonym

Activin-binding protein

Protein Construction

A DNA sequence encoding the mouse FST (P47931) (Met 1-Asn 317) was expressed with the Fc region of human IgG1 at the C-terminus.

Organism

Mouse

Expression Host

CHO Cells

QC Testing

Activity

Measured by its ability to neutralize Activin-mediated inhibition on MPC11 cell proliferation. The ED₅₀ for this effect is typically 40-200 ng/mL in the presence of 10 ng/mL rhActivin A.

Purity

> 90 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Gly 30

Molecular Mass

The recombinant mouse FST comprises 528 amino acids and has a predicted molecular mass of 58.4 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of the protein is approximately 66 kDa due to glycosylation.

Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Mouse Follistatin / FST (FS288) Protein (Fc Tag)
SDS-PAGE