

## General Information

### Gene Name Synonym

C-C Motif Chemokine 27, CC Chemokine ILC, Cutaneous T-Cell-Attracting Chemokine, CTACK, Eskine, IL-11 R-Alpha-Locus Chemokine, Skinkine, Small-Inducible Cytokine A27, CCL27, ILC, SCYA27

### Protein Construction

A DNA sequence encoding the Human CCL27/ILC/SCYA27 Phe25-Gly112 is expressed.

### Organism

Human

### Expression Host

E. coli

## QC Testing

### Purity

>95% as determined by SDS-PAGE.

### Endotoxin

<1.0 EU per  $\mu\text{g}$  protein as determined by the LAL method.

### Stability

Store for up to 12 months at  $-20^{\circ}\text{C}$  to  $-80^{\circ}\text{C}$  as lyophilized powder. Aliquots of reconstituted samples are stable at  $< -20^{\circ}\text{C}$  for 3 months.

## Formulation

Lyophilized from a  $0.2\ \mu\text{m}$  filtered solution of 20mM PB, 500mM NaCl, pH 7.5.

## Usage Guide


### Storage

Lyophilized protein should be stored at  $<-20^{\circ}\text{C}$ , though stable at room temperature for 3 weeks. Reconstituted protein solution can be stored at  $4-7^{\circ}\text{C}$  for 2-7 days.

### Reconstitution

Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than  $100\ \mu\text{g}/\text{ml}$ . Dissolve the lyophilized protein in ddH<sub>2</sub>O. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE



Human CCL27/ILC/SCYA27 recombinant protein