

MERS-CoV CoV Spike glycoprotein (aa 383-502, Fc Tag)



Catalog Number: 503542

General Information

Protein Construction

A DNA sequence encoding the spike protein fragment (Human betacoronavirus 2c EMC/2012)(AFS88936.1)(Cys383-Lys502) was fused with Fc region of mouse IgG at the C-terminus.

Organism

MERS-CoV

Expression Host

Baculovirus-Insect Cells

QC Testing

Activity

Measured by its binding ability in a functional ELISA.

Immobilized Spike (HCoV-EMC/2012) (Cat: 503542) at 10 µg/ml (100 µl/well) can bind biotinylated human ACE2-Fc (Cat: 501568) with a linear range of 0.03-4.0 µg/ml.

Purity

> 85 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per µg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Cys 383

Molecular Mass

The recombinant spike protein fragment (Human betacoronavirus 2c EMC/2012) comprises 354 amino acids and has a predicted molecular mass of 39.7 kDa. It migrates as an approximately 44 kDa band in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile 100mM Glycine, 10mM NaCl, pH 7.0.

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Novel coronavirus (HCoV-EMC/2012) Spike Protein fragment (aa 383-502, Fc Tag) SDS-PAGE