Human CXCL12 / SDF-1 beta (Fc Tag) recombinant protein

Catalog Number: 501323



General Information

Gene Name Synonym

C-X-C motif chemokine 12; Intercrine reduced in hepatomas; Pre-B cell growth-stimulating factor; SDF-1-beta(3-72); SDF-1-alpha(3-67)

Protein Construction

A DNA sequence encoding the N-terminally truncated human SDF1? (NP_000600.1) (Lys 22-Met 93) was expressed with the fused Fc region of human IgG1 at the N-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Measured by its ability to bind biotinylated recombinant human DPPIV in a functional ELISA.

Purity

> 95 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Glu 20

Molecular Mass

The recombinant human SDF1/Fc is a disulfide-linked homodimeric protein. The reduced monomer consists of 309 amino acids and predicts a molecular mass of 35.2 kDa. As a result of glycosylation, rhSDF1/Fc monomer is approximately 43 kDa in SDS-PAGE under reducing conditions.

Formulation

Lyophilized from sterile PBS, pH 7.4

- 1. 5 % trehalose and mannitol are added as protectants before lyophilization.
- 2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human CXCL12 / SDF1b Protein (Fc Tag) SDS-PAGE