## **Anti-PCGF2 antibody**

Catalog Number: 175763



#### Product name

Anti-PCGF2 antibody

#### **Specificity**

Human, Mouse, Rat, Chicken, Dog, Pig, Cow, Horse

#### Antibody description

Rabbit polyclonal antibody to PCGF2

## **Preparation**

This antigen of this antibody was klh conjugated synthetic peptide derived from human mel18/znf144 145-245

#### **Formulation**

Liquid, 0.01M TBS(pH7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

## **Storage**

Store at -20°C for one year. Avoid repeated freeze/thaw cycles. The lyophilized antibody is stable at room temperature for at least one month and for greater than a year when kept at -20°C. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4°C.

## **Clonality**

Polyclonal

#### Ig Type

Rabbit IgG

#### **Applications**

WB, IHC-P

#### **Dilutions**

WB:1:500-2000

IHC-P:1:400-800

#### **Validations**




Fig1: Tissue/cell: human colon carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;; Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum, C-0005) at 37°C for 20 min;; Incubation: Anti-Mel18/ZNF144 Polyclonal Antibody, Unconjugated 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010)




Fig2: Sample:; Muscle (Rat) Lysate at 40 ug; Primary: Anti- ZNF144 at 1/1000 dilution; Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution; Predicted band size: 38/60 kD;

Observed band size: 38 kD

# **Anti-PCGF2 antibody**

Catalog Number: 175763






Fig3: Paraformaldehyde-fixed, paraffin embedded (rat lung); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZNF144) Polyclonal Antibody, Unconjugated at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.