

Anti-LILRB2 antibody



Catalog Number: 175073

Product name

Anti-LILRB2 antibody

Specificity

Human

Antibody description

Mouse monoclonal antibody to LILRB2

Preparation

This antigen of this antibody was purified recombinant fragment of human *lilrb2* (aa: 51-184) expressed in *e. coli*.

Formulation

Liquid, 1*PBS with 0.05% sodium azide.

Storage

Store at +4°C after thawing. Aliquot store at -20°C. Avoid repeated freeze / thaw cycles.

Clonality

Monoclonal

Ig Type

Mouse IgG2a

Applications

WB, FC

Dilutions

WB: 1:500-1:2,000

FC: 1:100-1:200

Validations






Fig1: Western blot analysis of LILRB2 against human LILRB2 (AA: 51-184) recombinant protein. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Mouse IgG - HRP Secondary Antibody at 1:5,000 dilution was used for 1 hour




at room temperature.

Fig2: Western blot analysis of 175073# against HEK293 (1) and LILRB2 (AA: 51-184)-hIgGFc transfected HEK293 (2) cell lysate. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Mouse IgG - HRP Secondary Antibody at 1:5,000 dilution was used for 1 hour at room



temperature.

Fig3: Western blot analysis of 175073# against MOLT4 (1) and K562 (2) cell lysate. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody (1/500) was used in 5% BSA at room temperature for 2 hours. Goat Anti-Mouse IgG - HRP Secondary Antibody at 1:5,000 dilution was used for 1 hour at room



temperature.

Fig4: Flow cytometric analysis of LILRB2 was done on HL-60 cells. The cells were fixed, permeabilized and stained with the primary antibody (1/100) (green). After incubation of the primary antibody at room temperature for an hour, the cells were stained with a Alexa Fluor 488-conjugated goat anti-Mouse IgG Secondary antibody at 1/500 dilution for 30 minutes. Unlabelled sample was used as a control (cells without incubation with primary antibody; red).