## **Anti-BACE1 antibody**

Catalog Number: 106283



#### Product name

Anti-BACE1 antibody

#### **Immunogen**

Human BACE1 (His Tag) recombinant protein

## **Specificity**

Human BACE1

### **Antibody description**

Rabbit monoclonal to BACE1

### **Preparation**

This antibody was obtained from a rabbit immunized with purified, recombinant Human BACE1 (rh BACE1; NP\_036236.1; Met1-Thr457).

#### **Formulation**

0.2 µm filtered solution in PBS with 5% trehalose

## **Storage**

This antibody can be stored at  $2^{\circ}\text{C-8}^{\circ}\text{C}$  for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at  $-20^{\circ}\text{C}$  to  $-80^{\circ}\text{C}$ .

Preservative-Free.

Sodium azide is recommended to avoid contamination (final concentration 0.05%-0.1%). It is toxic to cells and should be disposed of properly. Avoid repeated freeze-thaw cycles.

## Clonality

Monoclonal

#### Ig Type

Rabbit IqG

## **Applications**


FCM, IF, ICC/IF

#### **Dilutions**

FCM: 0.5-2 µg/Test


ICC/IF:  $10-25 \mu g/mL$ 

#### **Validations**



BACE1 Antibody, Rabbit MAb, Flow cytometric analysis

Flow cytometric analysis of Human BACE1 expression on Jurkat cells. The cells were treated according to manufacturer's manual (BD Pharmingen™ Cat. No. 554714), stained with purified anti-Human BACE1, then a FITC-conjugated second step antibody. The fluorescence histograms were derived from gated events with the forward and side light-scatter characteristics of intact cells.



BACE1 Antibody, Rabbit MAb, Immunofluorescence

Immunofluorescence staining of Human BACE1 in Hela cells. Cells were fixed with 4% PFA,

# **Anti-BACE1 antibody**

Catalog Number: 106283



permeabilzed with 1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human AK4 monoclonal antibody (15  $\mu$ g/ml) at 37°C 1 hour. Then cells were stained

with the Alexa Fluor® 488-conjugated Goat Antirabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to cytoplasm.