

General Information

Gene Name Synonym

V-set and immunoglobulin domain containing 4

Protein Construction

A DNA sequence encoding the extracellular domain (Met 1-Pro 187) of mouse VSIG4 (NP_808457.1) precursor was fused with the C-terminal polyhistidine-tagged Fc region of human IgG1 at the C-terminus.

Organism

Mouse

Expression Host

Human Cells

QC Testing

Purity

> 95 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

His 20

Molecular Mass

The recombinant mouse VSIG4/Fc is a disulfide-

linked homodimer after removal of the signal peptide. The reduced monomer consists of 416 amino acids and has a predicted molecular mass of 47 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of rmVSIG4/Fc monomer is approximately 60 kDa due to glycosylation.

Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C . It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Mouse VSIG4 Protein (His & Fc Tag) SDS-PAGE