

Human CD34 (Fc Tag) recombinant protein



Catalog Number: 500814

General Information

Protein Construction

A DNA sequence encoding the extracellular domain of human CD34 precursor (NP_001020280.1) (Met 1-Thr 290) was fused with the Fc region of human IgG1 at the C-terminus.

Organism

Human

Expression Host

Human Cells

QC Testing

Activity

Measured by the ability of the immobilized protein to support the adhesion of the HUVEC human umbilical vein endothelial cell line. When 4×10^4 cells/well are added to human CD34 coated plates (0.8 $\mu\text{g/ml}$, 100 $\mu\text{l/well}$), approximately >40 % will adhere after one hour at 37 °C.

Purity

> (67.1+28.7) % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Ser 32

Molecular Mass

The recombinant human CD34/Fc is a disulfide-linked homodimer generated after removal of the signal peptide. The reduced monomer consists of 497 amino acids and has a predicted molecular mass of 54 kDa. In SDS-PAGE under reducing conditions, the apparent molecular mass of the protein is 116 and 96 kDa due to different glycosylation.

Formulation

Lyophilized from sterile PBS, pH 7.4

1. 5 % trehalose and mannitol are added as protectants before lyophilization.

2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Human CD34 Protein (Fc Tag) SDS-PAGE