

Mouse Cystatin 7 / CST7 (Fc Tag) recombinant protein



Catalog Number: 501334

General Information

Gene Name Synonym

Cystatin-7; Cystatin-like metastasis-associated protein; Leukocystatin

Protein Construction

A DNA sequence encoding the mouse CST7 (O89098) (Met1-Gln144) was expressed, fused with the Fc region of human IgG1 at the C-terminus.

Organism

Mouse

Expression Host

Human Cells

QC Testing

Activity

Measured by its ability to inhibit active Cathepsin L cleavage of a fluorogenic peptide substrate Z-LR-AMC, R&D Systems, Catalog # ES008. The IC50 is < 6 nM.

Purity

> 93 % as determined by SDS-PAGE

Endotoxin

< 1.0 EU per μg of the protein as determined by the LAL method

Stability

Samples are stable for up to twelve months from date of receipt at -70°C

Predicted N terminal

Ala 19

Molecular Mass

The recombinant mouse CST7 /Fc is a disulfide-linked homodimer. The reduced monomer comprises 367 amino acids and has a predicted molecular mass of 41.4 KDa. The apparent molecular mass of the protein is approximately 42 KDa in SDS-PAGE under reducing conditions due to glycosylation.

Formulation

Lyophilized from sterile 25mM Tris, 0.15M NaCl, pH 7.5

1. 5 % trehalose and mannitol are added as protectants before lyophilization.
2. Please contact us for any concerns or special requirements.

Usage Guide


Storage

Store it under sterile conditions at -20°C to -80°C . It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

Reconstitution

Adding sterile water, prepare a stock solution of 0.25 mg/ml. Concentration is measured by UV-Vis.

SDS-PAGE



Mouse Cystatin 7 / CST7 Protein (Fc Tag) SDS-PAGE